

Improvement in Latent Fingerprint Detection on Thermal Paper using 5,6-Dimethoxy-1,2-indandione/PVP

Yabin Zhao, Yaxian Feng and Yaping Luo*

Department of Forensic Science, Chinese People's Public Security University, Beijing, China

*Corresponding author: Yaping Luo, Department of Forensic Science, Chinese People's Public Security University, Beijing 100038, China, Fax: +86 010-83903380, Tel: +86 13501265813, E-mail: lyp6698@163.com

Citation: Yabin Zhao, Yaxian Feng, Yaping Luo (2013) Improvement in Latent Fingerprint Detection on Thermal Paper using 5,6-Dimethoxy-1,2-indandione/PVP. J Forensic Sci Criminol 1(2): 202. doi: 10.15744/2348-9804.1.202

Received Date: September 24, 2013 Accepted Date: October 15, 2013 Published Date: October 28, 2013

Abstract

A new method for improvement of 1,2-indandione-treated latent fingerprints on thermal paper will be described in this paper. Treatment with conventional techniques like ninhydrin in petroleum ether or DFO solution produces a black color on the thermosensitive side of thermal paper. The new method using 5,6-dimethoxy-1,2-indandione with polyvinyl pyrrolidone (PVP) can reduce this dark staining without removing the thermosensitive layer and parts of the developed fingerprints. An advantage of this method is that the developed fluorescent fingerprints appear in sharp lines and high contrast. The developed fingerprints can be observed when excited in the 535 nm region and using an orange-red goggles. There is no background coloration in this method. In addition, some tests were performed, leading to an optimized working solution, which charges the paper with a minimum of chemicals, is cheap, and enables a large quantity of papers to be treated in a short time and without involving any pre- or post treatment.

Keywords: 5,6-Dimethoxy-1,2-indandione; Polyvinylpyrrolidones; Thermal paper; Latent fingerprint

Introduction

The thermal paper has been extensively used in modern day life, such as fax machines, ATM receipts, store receipts, lotteries, bus tickets etc. For this reason, it is collected as a physical evidence from the crime scene more and more common. However, most thermosensitive surfaces of thermal paper turn black when they come into contact with polar organic solvents or are exposed to high temperature. This black background staining reduces the contrast of the developed fingerprints, often rendering them useless for identification purposes [1]. So, the continuing need for new reagents and methods to develop and visualize latent fingerprints on thermal papers for forensic examination has been well-investigated by forensic science researchers. A desirable attribute of fingerprint development techniques is high sensitivity, selectivity, and maximization of the contrast between the developed fingerprints and the background.

In 1991, Takatsu et al. [2] showed that a solution of alkyl analogues of ninhydrin could be used to develop latent fingerprints on thermal paper, but only obtained partial development of the fingerprints [2,3]. Using INON (2-hydroxy-2-(3,5,5-trimethyl-hexyloxy)-indan-1,3-dione) they reduced the blackening of the surface and obtained the better results [2]. By fuming thermal papers with dimethylaminocinnamaldehyde (DMAC) and iodine, Brennan and Jasuja were able to develop clear fingerprints

with no background coloration [4,5], although vapor-phase staining method had been known for many years as a general development technique for depositions on paper. To enhance ninhydrin treated latent fingerprints on thermal paper, Schwarz used a solution containing pyrrolidone based compounds referred to as 'G3', to decolorize the blackened paper [6]. The majority of the techniques discussed above suffer from one or more drawbacks such as being complex or cumbersome, some requiring pre- or post treatment, or being less efficient at developing aged fingerprints. In 2010, Schwarz improved the methodology by adding polyvinyl pyrrolidone (PVP) into the solution of ninhydrin and achieved very promising results which there is no black background anymore. No mark of fluorescence has been noted by the purple fingerprint developed by ninhydrin. There will be background interference when the developed purple mark was on the printed words [7]. This encouraged us to pursue a method for the development of latent fingerprints on thermal paper solving the problems. Quite recently, we have reported a new DFO/PVP development method that can be successfully applied to develop fresh as well as aged latent fingerprints on thermal paper with no background coloration [8]. 1,2-Indanedione belongs to a class of compounds which have demonstrated great potential in the processing of latent fingerprints. It has been proved to be a viable alternative for the detection of latent fingerprints on porous surfaces, with more fingerprints being developed using this reagent on real samples than both DFO, ninhydrin and a combination of the two reagents. Although the synthesis of

1,2-indanediones had been reported many years ago, these compounds had never been tested before on latent prints until 1997. 1,2-Indanedione was proposed as a fingerprint reagent by Ramotowski [9]. Following this discovery, much effort has been devoted to the latent fingerprints development by using 1,2-indanediones. Researchers worldwide have conducted research into the optimization and evaluation of 1,2-indanedione as an adequate reagent for fingerprint detection [10-15].

An evaluation of 5,6-dimethoxy-1,2-indandione in latent fingerprints detection began with the work of Almog, et al. [11]. The fluorescence of this new compound was comparable to the best ninhydrin analogs. The fluorescence was significantly enhanced by the subsequent application of a zinc nitrate solution. We envisioned that using the 5,6-dimethoxy-1,2-indandione/PVP system, where the presence of a latent fingerprint was visualized by luminescence, would provide a simple and effective method for latent fingerprint detection on thermal paper. In the current work, the mixture solution of 5,6-dimethoxy-1,2-indandione with PVP was utilized as working solution to detect latent fingerprints on thermal paper. Integrating PVP into a 5,6-dimethoxy-1,2-indandione solution prevents the black staining, and the developed fingerprints appear in clear contrast to the background under the fluorescence conditions.

Using the resultant optimized formulation, performance comparisons were then made. The quality score of developed marks was used to evaluate the effects. The sensitivity, selectivity and stability of the method were also investigated.

Material and methods

Materials

All the solvents (AR grade) were purchased from Beijing Chemical Co., Ltd (Beijing) and used without further purification. 5,6-Dimethoxy-1,2-indandione was supplied by Chinese People's Public Security University. PVP with lower molar mass Kolloidon® 12 PF (Mw = 2000–3000; BASF ChemTrade GmbH, Burgbernheim, Germany) was employed.

Preparation of the 5,6-dimethoxy-1,2-indandione Solution consisted of fully dissolving 0.8 g of 5,6-dimethoxy-1,2-indandione in 90 ml ethyl acetate and 10 ml acetic acid. Then mixing thoroughly with 80ml zinc chloride solution (prepared with zinc chloride, ethyl alcohol, petroleum ether and ethyl acetate) and 820 ml petroleum ether.

The DFO/PVP working solution: The best ratio of DFO solution in PVP(5%), in ethanol solution, was 1:10 (v/v). Fingerprints developed on thermal paper by DFO/PVP were visualized at 515 nm excitation and observed with an orange-red goggles.

Fingerprint samples

5 individuals (1 female, 4 males) aged from 20-25 were chosen as donors for the experiments. Donors were requested not to wash their hands during the experiments in order to produce

natural fingerprints. Donors were asked to deposit a depletion series, which consisted of 5 successive contacts on the thermal paper with a single finger.

Every individual followed and repeated the same procedure for 10 times. We tried our best to ensure that each volunteer impress their fingerprints with similar force and within the same duration. The fingerprints were stored under room temperature in the laboratory for 1, 3, 5, 7, 15 and up to 30 days before being treated. The comparison between different methods was performed by cutting the series of fingerprints in halves before dipping them in their respective staining solutions. Developed marks were photographed using a digital camera (Nikon D70, AF Micro-Nikkor 60mm f / 2.8D, Japan).

Detection of latent fingerprints

The detection of the latent fingerprints was performed as follows: (1) Each sample was immersed in various staining solutions for 10s; (2) After that, the treated samples were heated in an oven at 100 °C, relative humidity 60% for 10 min. (3) The samples were then observed in the luminescence mode using a Polilight lamp with the emission set at an excitation wavelength of 535 nm.

Results and discussions

Optimization of 5,6-dimethoxy-1,2-indandione/PVP staining solution

PVP is a type of non-ionic polymer, with low volatility and nontoxic, it can be easily dissolved in polar solution and difficult to dissolve in the less or non-polar solvents. To increase the solubility and fluorescence, the influence of solvent and the ratio of PVP to 5,6-dimethoxy-1,2-indandione on detecting latent fingerprints was also explored. It was determined that use of ethanol as a solvent to assist solubility exerts a substantial influence on the sensitivity of the reagent. The optimum ratio of 5,6-dimethoxy-1,2-indandione solution with PVP, in ethanol solution, was 1:7.5(v/v) and the best concentration of PVP in the ethanol solution was determined to be around 5% (Table 1). Encouraged by this result, further optimization studies using fingerprints deposited on thermal papers were extended to different remaining days and afforded similar results.

Concentration of PVP solution	PVP to indandione ratio				
	1: 1	1: 2.5	1: 5	1: 7.5	1: 10
1%	+	+	++	+++	++
2.5%	+	+	++	+++	++
5%	+	+	+++	++++	+++
7.5%	+	++	+++	++++	+++
10%	+	+	++	+++	++

The test result is indicated by +. +++++ indicates very high quality fingerprints, +++ indicates clear visible fingerprints, ++ indicates that the fingerprints are clear in general but certain parts are difficult to identify, + indicates that the fingerprints are incomplete, not clear enough for identification; - indicates that the fingerprints do not appear on thermal paper.

Table 1: Detection results developed with 5,6-dimethoxy-1,2-indandione/PVP for fingerprints impressed on thermal paper within 1 day.

Efficiency of the new PVP/DFO staining solution

Figure 1 shows a typical example of freshly deposited latent "natural" fingerprint, developed by 5,6-dimethoxy-1,2-indandione/PVP under the above mentioned conditions. Fresh fingerprints developed with 5,6-dimethoxy-1,2-indandione/PVP solution for all donors appear in sharp lines, high contrast, excellent fluorescence and without background coloration. The fluorescent fingerprints can be observed and photographed when excited in the 535 nm region through an orange-red goggles.




Figure 1: Photographic images of a latent "natural" fingerprint from a single donor deposited on thermal paper. The fresh latent fingerprint developed with 5,6-dimethoxy-1,2-indandione/PVP solution, visualised at 535 nm excitation and observed with an orange-red goggles.

Comparison of the 5,6-dimethoxy-1,2-indandione/PVP and the, 6-dimethoxy-1,2-indandione

The comparison between 5,6-dimethoxy-1,2-indandione/PVP formulation with those obtained only using 5,6-dimethoxy-1,2-indandione was performed by splitting the fingerprints in half before dipping them in their respective staining solutions (5,6-dimethoxy-1,2-indandione/PVP for the left halves and 5,6-dimethoxy-1,2-indandione for the right halves). Latent fingerprints on thermal papers developed with 5,6-dimethoxy-1,2-indandione turned dark on the thermosensitive side, while those developed with 5,6-dimethoxy-1,2-indandione/PVP showed stronger luminescence, excellent quality of ridge detail and better contrast with the background. (Figure 2).




Figure 2: Latent fingerprint developed on thermal paper by (A) 5,6-Dimethoxy-1,2-indandione/PVP (left halve) and (B) 5,6-Dimethoxy-1,2-indandione (right halve). All the developed fingerprints were visualized at 535 nm excitation, observed using orange-red goggles. The results from all samples test showed that the change trend of enhancement for different donors were consistent.

Comparison of the 5,6-dimethoxy-1,2-indandione/PVP and the PVP/DFO

Subsequently, we compared the efficiency of DFO/PVP with 5,6-dimethoxy-1,2-indandione/PVP methods. The merits of the new method are clear. Figure 3 shows an example where both methods detected fingerprints on the thermal paper without turning dark on the thermosensitive side, while those developed with 5,6-dimethoxy-1,2-indandione/PVP provided greater fluorescence, better detection sensitivity and clearer detail than that obtained with PVP/DFO.



Figure 3: Latent fingerprint developed on thermal paper with (A) 5,6-Dimethoxy-1,2-indandione/PVP, polilight lamp at 535 nm excitation, observed using orange-red goggles in the luminescence mode. (B) One-step PVP/DFO, polilight lamp at 515 nm excitation, observed using orange-red goggles in the luminescence mode. The results from all samples test showed that the change trend of enhancement for different donors were consistent.

Sensitivity and stability of the 5,6-dimethoxy-1,2-indandione/PVP treatment for latent fingerprints

Next, the sensitivity and stability of the one-step 5,6-dimethoxy-1,2-indandione/PVP treatment for latent fingerprints were also explored. A depletion series of marks with decreasing quantities of material deposited were test. The typical depletion series consisting of five consecutive finger depositions (Figure 4) showed that the new method provided sharp and clear development of latent fingerprints, without background staining, even dramatically diminished the amounts of residue deposited in fingerprints, the image still showing ridge detail (Figure 4E). Furthermore, the efficiency of development for all samples showed that principally fingerprints from different donors were consistently visualized and the tendency of the depletion for different donors were similar.

Conclusions

In this report, a new one-step indandione /PVP development method has been proposed for detecting latent fingerprints on thermal paper with no background coloration. 5,6-Dimethoxy-1,2-indandione/PVP has been successfully synthesized and we suggest its use as the working solution to detect latent fingerprints on thermal papers. The new technique could obtain even better results compared with the existing efficient reagents. It is a simple method to detect latent fingerprints that provides sharp and clear development of latent fingerprints, without background staining, dramatically diminished the



Figure 4: Illustration of typical depletion series consisting of five consecutive finger depositions on thermal paper, aged for 1 day and treated with the 5,6-dimethoxy-1,2-indandione/PVP. All the developed fingermarks were visualised at 535 nm excitation and observed using an orange-red goggles.




Figure 5: Photographs of 5,6-dimethoxy-1,2-indandione/PVP developed latent fingermarks on thermal papers aged for 1 day, 3 days, 5 days, 7 days, 15 days and 30 days.

steps, with simple operation. This technique establishes a new possibility for the detection of latent fingermarks in forensic science. Finally, we have checked the shelf-life of synthesized developer, the visual contrast of developed fingermarks did not change with the developer stored time over periods of several months and the developer also did not deteriorate significantly when stored under ambient conditions.

Acknowledgements

The authors would like to thank the programs of Beijing Municipal commission of education for the grant provided to support this research .

References

- Schwarz L (2003) Trace evidence on thermal paper. *Criminology* 57: 246-9
- Takatsu M, Kageyama H, Hirata K, Akashi S, Yoko T T, et al. (1991) Development of a new method to detect latent fingerprint on thermal paper with o-Alkyl derivative of ninhydrin. *Kagaku Keisatsu Kenkyusho Hokoku* 44: 1-6.
- Takatsu M, Akaishi S, Kageyama H (1992) Fingerprint detection reagents and the preparation thereof. *Japanese Patent* 4-124156: 1-8.
- Brennan JS, Bramble SK, Crabtree S, Wright G (1995) Fuming of latent fingermarks using dimethylaminocinnamaldehyde. *J Forensic Ident* 45: 373-80.
- Jasuja OP, Singh G (2009) Development of latent fingermarks on thermal paper: preliminary investigation into use of iodine fuming. *Forensic Sci Int* 192: e11-16.
- Schwarz L, Klenke I (2007) Enhancement of ninhydrin- or DFO-treated latent fingerprints on thermal paper. *J Forensic Sci* 52: 649-55.
- Schwarz L, Klenke I (2010) Improvement in latent fingerprint detection on thermal paper using a one-step ninhydrin treatment with polyvinylpyrrolidones (PVP). *J Forensic Sci* 55: 1076-79.
- Luo YP, Zhao YB, Liu S (2013) Evaluation of DFO/PVP and its application to latent fingermarks development on thermal paper. *Forensic Sci Int* 229: 75-9.
- Ramotowski R, Cantu AA, Joulie MM, Petrovskaia O (1997) 1,2-Indanediones: a preliminary evaluation of a new class of amino acid visualizing compounds. *Fingerprint Whorld* 23: 131-40.
- Roux C, Jones N, Lennard C, Stoilovic M (2000) Evaluation of 1,2-indanedione and 5,6-dimethoxy-1,2-indanedione for the detection of latent fingerprints on porous surfaces. *J Forensic Sci* 45: 761-9.
- Almog J, Springer E, Wiesner S, Frank A, Khodzhaev O, et al. (1999) Latent fingerprint visualization by 1,2-indanedione and related compounds: preliminary results. *J Forensic Sci* 44: 114-40.
- Wiesner S, Springer E, Sasson Y, Almog J (2001) Chemical development of latent fingerprints: 1,2-indanedione has come of age. *J Forensic Sci* 46: 1082-84.
- Hauze D, Petrovskaia O, Taylor B, Joulie MM, Ramotowski R, et al. (1998) 1,2-Indanediones: new reagents for visualizing the amino acid components of latent prints. *J Forensic Sci* 43: 744-7.
- Kasper S, Minnillo D, Rockhold A (2002) Validating IND (1,2-indanedione). *Forensic Sci Commun* 4.
- Merrick S, Gardner S, Sears VG, Hewlett DF (2002) An operational trial of ozone friendly DFO and 1,2-indanedione formulations for latent fingerprint detection. *J Forensic Ident* 52: 595-605.

Submit your next manuscript to Annex Publishers and benefit from:

- ▶ Easy online submission process
- ▶ Rapid peer review process
- ▶ Online article availability soon after acceptance for Publication
- ▶ Open access: articles available free online
- ▶ More accessibility of the articles to the readers/researchers within the field
- ▶ Better discount on subsequent article submission

Submit your manuscript at

<http://www.annexpublishers.com/paper-submission.php>